

provided in #15  
U.S.S.N. 08/488,087  
Examiner Ford

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office

March 17, 1998

RECEIVED


JUL 16 1998

GROUP 180

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 07/473,318

FILING DATE: February 1, 1990



By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS

D. SMITH

Certifying Officer


BEST AVAILABLE COPY

PATENT APPLICATION SERIAL NO. **07 473318**

U.S. DEPARTMENT OF COMMERCE  
PATENT AND TRADEMARK OFFICE  
FEE REC'D SHEET

140 02/16/90 07473318

1 201 1,163.00 CK



EXPRESS MAIL NO. B011 781 05H

07 473318

0510.013

TO ALL WHOM IT MAY CONCERN:

Be it known that I, Dennis C. Liotta, a citizen of the United States of America, and Woo-Baeg Choi, a citizen of the Republic of Korea, residing at 793 Post Road Way, Stone Mountain, Georgia 30088; and 3215A Flowers Road, S., Atlanta, Georgia 30341, respectively, have invented new and useful improvements in  
a

5c1  
METHOD AND COMPOSITIONS FOR THE SYNTHESIS OF DCU-189  
AND RELATED COMPOUNDS

for which the following is a specification.

4/8

\$ 1163.00 - 101- R

07 473318



ACKNOWLEDGEMENT

The invention described herein was made with Government support under grant no. 5-21935 awarded by the National Institutes of Health. The Government has certain rights in this invention.

BACKGROUND OF THE INVENTION

The present invention relates to methods and compositions for preparing antiviral nucleoside analogs, particularly BCH-189 (2',3'-dideoxy-3'-thia-cytidine). More particularly, the invention relates to the selective synthesis of the  $\beta$ -isomer of BCH-189 and related compounds as well as the selective synthesis of enantiomerically-enriched BCH-189 and related compounds.

In 1981, documentation began on the disease that became known as Acquired Immune Deficiency Syndrome (AIDS), as well as its forerunner AIDS Related Complex (ARC). In 1983, the cause of the disease AIDS was established as a virus named the Human Immunodeficiency Virus type 1 (HIV-1). Usually, a person infected with the virus will eventually develop AIDS; in all known cases of AIDS the final outcome has always been death.

The disease AIDS is the end result of an HIV-1 virus following its own complex life cycle. The virion life cycle begins with the virion attaching itself to the host human T-4 lymphocyte immune cell through the bonding of a glycoprotein on the surface of the virion's protective coat with the CD4 glycoprotein on the lymphocyte cell. Once attached, the virion sheds its glycoprotein coat, penetrates into the membrane of the host cell, and uncoats its RNA. The virion enzyme, reverse transcriptase, directs the process of transcribing the RNA into single stranded DNA. The viral RNA is degraded and a second DNA strand is created. The now double-stranded DNA is integrated into the human cell's genes and those genes are used for cell reproduction.

At this point, the human cell carries out its reproductive process by using its own RNA polymerase to transcribe the integrated DNA into viral RNA. The viral RNA is translated into glycoproteins, structural proteins, and viral enzymes, which assemble with the viral RNA intact. When the host cell finishes the reproductive step, a new virion cell, not a T-4 lymphocyte, buds forth. The number of HIV-1 virus cells thus grows while the number of T-4 lymphocytes decline.

The typical human immune system response, killing the invading virion, is taxed because a large portion of the virion's life cycle is spent in a latent state within the immune cell. In

addition, viral reverse transcriptase, the enzyme used in making a new virion cell, is not very specific, and causes transcription mistakes that result in continually changed glycoproteins on the surface of the viral protective coat. This lack of specificity decreases the immune system's effectiveness because antibodies specifically produced against one glycoprotein may be useless against another, hence reducing the number of antibodies available to fight the virus. The virus continues to grow while the immune response system continues to weaken. Eventually, the HIV largely holds free reign over the body's immune system, allowing opportunistic infections to set in and ensuring that, without the administration of antiviral agents and/or immunomodulators, death will result.

There are three critical points in the virus's life cycle which have been identified as targets for antiviral drugs: (1) the initial attachment of the virion to the T-4 lymphocyte, or macrophage, site, (2) the transcription of viral RNA to viral DNA, and (3) the assemblage of the new virion cell during reproduction.

Inhibition of the virus at the second critical point, the viral RNA to viral DNA transcription process, has provided the bulk of the therapies used in treating AIDS. This transcription must occur for the virion to reproduce because the virion's genes are encoded in RNA; the host cell reads only DNA.

By introducing drugs that block the reverse transcriptase from completing the formation of viral DNA, HIV-1 replication can be stopped.

Nucleoside analogs, such as 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxycytidine (DDC), 2',3'-dideoxythymidinene (D4T), 2',3'-dideoxyinosine (DDI), and various fluoro-derivatives of these nucleosides are relatively effective in halting HIV replication at the reverse transcriptase stage. Another promising reverse transcriptase inhibitor is 2',3'-dideoxy-3'-thia-cytidine (BCH-189), which contains an oxathiolane ring substituting for the sugar moiety in the nucleoside.

AZT is a successful anti-HIV drug because it sabotages the formation of viral DNA inside the host T-4 lymphocyte cell. When AZT enters the cell, cellular kinases activate AZT by phosphorylation to AZT triphosphate. AZT triphosphate then competes with natural thymidine nucleosides for the receptor site of HIV reverse transcriptase enzyme. The natural nucleoside possesses two reactive ends, the first for attachment to the previous nucleoside and the second for linking to the next nucleoside. The AZT molecule has only the first reactive end; once inside the HIV enzyme site, the AZT azide group terminates viral DNA formation because the azide cannot make the 3',5'-phosphodiester with the ribose moiety of the following nucleoside.

AZT's clinical benefits include increased longevity, reduced frequency and severity of opportunistic infections, and increased peripheral CD4 lymphocyte count. Immunosorbent assays for viral p24, an antigen used to track HIV-1 activity, show a significant decrease with use of AZT. However, AZT's benefits must be weighed against the severe adverse reactions of bone marrow suppression, nausea, myalgia, insomnia, severe headaches, anemia, peripheral neuropathy, and seizures. Furthermore, these adverse side effects occur immediately after treatment begins whereas a minimum of six weeks of therapy is necessary to realize AZT's benefits.

Both DDC and D4T are potent inhibitors of HIV replication with activities comparable (D4T) or superior (DDC) to AZT. However, both DDC and D4T are converted to their 5' triphosphates less efficiently than their natural analogs and are resistant to deaminases and phosphorylases. Clinically, both compounds are toxic. Currently, DDI is used in conjunction with AZT to treat AIDS. However, DDI's side effects include sporadic pancreatitis and peripheral neuropathy. Initial tests on 3'-fluoro-2'-3'-dideoxythymidine show that its anti-viral activity is comparable to that of AZT.

Recent tests on BCH-189 have shown that it possesses anti-HIV activity similar to AZT and DDC, but without the cell toxicity which causes the debilitating side effects of AZT and



DDC. A sufficient quantity of BCH-189 is needed to allow clinical testing and treatment using the drug.

The commonly-used chemical approaches for synthesizing nucleosides or nucleoside analogs can be classified into two broad categories: (1) those which modify intact nucleosides by altering the carbohydrate, the base, or both and (2) those which modify carbohydrates and incorporate the base, or its synthetic precursor, at a suitable stage in the synthesis. Because BCH-189 substitutes a sulfur atom for a carbon atom in the carbohydrate ring, the second approach is more feasible. The most important factor in this latter strategy involves delivering the base from the  $\beta$ -face of the carbohydrate ring in the glycosylation reaction because only the  $\beta$ -isomers exhibit useful biological activity.

It is well known in the art that the stereoselective introduction of bases to the anomeric centers of carbohydrates can be controlled by capitalizing on the neighboring group participation of a 2-substituent on the carbohydrate ring (Chem. Ber. 114:1234 (1981)). However, BCH-189 and its analogs do not possess a 2-substituent and, therefore, cannot utilize this procedure unless additional steps to introduce a functional group that is both directing and disposable are incorporated into the synthesis. These added steps would lower the overall efficiency of the synthesis.

It is also well known in the art that "considerable amounts of the undesired  $\alpha$ -nucleosides are always formed during the synthesis of 2'-deoxyribosides" (Chem. Ber. 114:1234, 1244 (1981)). Furthermore, this reference teaches that the use of simple Friedel-Crafts catalysts like  $\text{SnCl}_4$  in nucleoside syntheses produces undesirable emulsions upon the workup of the reaction mixture, generates complex mixtures of the  $\alpha$  and  $\beta$ -isomers, and leads to stable  $\sigma$ -complexes between the  $\text{SnCl}_4$  and the more basic silyated heterocycles such as silyated cytosine. These complexes lead to longer reaction times, lower yields, and production of the undesired unnatural N-3-nucleosides. Thus, the prior art teaches the use of trimethylsilyl triflate or trimethylsilyl perchlorate as a catalyst during the coupling of pyrimidine bases with a carbohydrate ring to achieve high yields of the biologically active  $\beta$ -isomers. However, the use of these catalysts to synthesize BCH-189 or BCH-189 analogs does not produce the  $\beta$ -isomer preferentially; these reactions result in approximately a 50:50 ratio of the isomers.

Thus, there exists a need for an efficient synthetic route to BCH-189 and its analogs. There also exists a need for a stereoselective synthetic route to the biologically active isomer of these compounds,  $\beta$ -BCH-189 and related  $\beta$ -analogues. Furthermore, there exists a need for a stereoselective synthetic route to enantiomerically-enriched  $\beta$ -BCH-189 because the other enantiomer is inactive and, therefore, represents a 50% impurity.

### SUMMARY OF THE INVENTION

The present invention relates to the discovery of a surprisingly efficient synthetic route to BCH-189 and various analogs of BCH-189 from inexpensive precursors with the option of introducing functionality as needed. This synthetic route allows the stereoselective preparation of the biologically active isomer of these compounds,  $\beta$ -BCH-189 and related compounds.

Furthermore, the stereochemistry at the nucleoside 4' position can be controlled to produce enantiomerically-enriched  $\beta$ -BCH-189 and its analogs.

The term "BCH-189 analogs" is meant to refer to nucleosides that are formed from pyrimidine bases substituted at the 5 position that are coupled to substituted 1,3-oxathiolanes.

The method of the present invention includes ozonizing an allyl ether or ester having the formula  $\text{CH}_2=\text{CH}-\text{CH}_2-\text{OR}$ , in which R is a protecting group, such as an alkyl, silyl, or acyl group, to form a glycoaldehyde having the formula  $\text{OHC}-\text{CH}_2-\text{OR}$ ; adding thioglycolic acid to the glycoaldehyde to form a lactone of the formula 2-(R-oxy)-methyl-5-oxo-1,3-oxathiolane; converting the lactone to its corresponding carboxylate at the 5 position of the oxathiolane ring; coupling the acetate with a silylated pyrimidine base in the presence of  $\text{SnCl}_4$  to form the  $\beta$ -isomer of a 5'-(R-oxy)-2',3'-dideoxy-3'-thia- nucleoside analog; and

replacing the R protecting group with a hydrogen to form BCH-189 or an analog of BCH-189.


The invention can be used to produce BCH-189 or BCH-189 analogs that are enantiomerically-enriched at the 4' position by selecting an appropriate R protecting group to allow stereoselective selection by an enzyme. For instance, the R protecting group can be chosen such that the substituent at the 2 position of the oxathiolane lactone is butyryloxy to permit stereoselective enzymatic hydrolysis by pig liver esterase. The resulting optically active hydrolyzed lactone can then be converted to its corresponding diacetate and coupled with a silylated pyrimidine base as above.

Accordingly, one of the objectives of this invention is to provide an efficient method for preparing the  $\beta$ -isomer of BCH-189 and analogs of BCH-189 in high yields. Furthermore, it is an objective of this invention to provide a synthetic method to produce only one optical isomer, rather than a racemic mixture, of BCH-189 and analogs of BCH-189. A further object of this invention is to provide a synthetic route to produce  $\beta$ -BCH-189 that is enantiomerically-enriched.

Additionally, an objective of this invention is to provide intermediates from which BCH-189 or BCH-189 analogs can be synthesized of the formula 2-(R-oxymethyl)-5-acyloxy-1,3-


oxathiolane, wherein R is a protecting group, such as alkyl, silyl, or acyl, and a method of preparing these compounds. Furthermore, it is an object of this invention to provide enantiomerically-enriched 2-acetoxymethyl-5-acetoxy-1,3-oxathiolane and 2-butoxymethyl-5-oxo-1,3-oxathiolane and methods of preparing these compounds.

Another objective of this invention is to provide intermediates from which BCH-189 or BCH-189 analogs can be synthesized of the formula:



wherein R is a protecting group, such as alkyl, silyl, or acyl, and Y can be hydrogen, methyl, halo, alkyl, alkenyl, alkynyl, hydroxalkyl, carboxalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl, and methods of preparing these compounds.

Furthermore, this invention provides intermediates from which BCH-189 or BCH-189 analogs can be synthesized of the formula:



wherein R is a protecting group, such as alkyl, silyl, or acyl, and Y can be hydrogen, methyl, halo, alkyl, alkenyl, alkynyl, hydroxalkyl, carboxalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl, and methods of preparing these compounds.

#### BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates one embodiment of a synthesis of BCH-189 and BCH-189 analogs according to the present invention;


Figure 2 illustrates one embodiment of the synthesis of BCH-189 according to the present invention;

Figure 3 illustrates one embodiment of the synthesis of 5-methylcytidine and thymidine derivatives of BCH-189 according to the present invention; and

Figure 4 illustrates one embodiment of the synthesis of enantiomerically-enriched BCH-189 according to the present invention.

### DETAILED DESCRIPTION OF THE INVENTION

BCH-189 is a compound of the formula:




The process of the present invention for preparing BCH-189 and BCH-189 analogs is set forth in Fig. 1. An allyl ether or ester 1 is ozonized to give an aldehyde 2, which reacts with thioglycolic acid to give a lactone 3. The lactone 3 is treated with a reducing agent, followed by a carboxylic anhydride, to produce the carboxylate 4. This carboxylate is coupled with a silylated pyrimidine base in the presence of a Lewis acid that can catalyze stereospecific coupling, such as  $\text{SnCl}_4$ , to yield the  $\beta$ -isomer of the substituted nucleoside 5 in essentially a 100:0 ratio of  $\beta$ : $\alpha$  isomers. The substituted nucleoside 5 is deprotected to produce BCH-189 or BCH-189 analog 6.

This procedure can be tailored to produce BCH-189 or BCH-189 analogs that are enantiomerically-enriched at the 4' position by selecting an appropriate R protecting group to allow stereoselective enzymatic hydrolysis of 1 by an enzyme such as pig liver esterase, porcine pancreatic lipase, or subtilisin or other enzymes that hydrolyze 1 in a stereoselective fashion. The resulting optically active 1 can be converted to

enantiomerically-enriched carboxylate 4 and coupled with a silylated pyrimidine base as above to produce enantiomerically-enriched BCH-189 or BCH-189 analogs.

The protecting group R in 1 can be selected to provide protection for the corresponding alcohol until the final step in the synthesis is carried out (deprotection of 5 to form 6). Additionally, the protecting group can be selected, if desired, to provide an additional recognition site for an enzyme to be used later in an enantio-selective hydrolysis reaction. Any group that functions in this manner may be used. For instance, alkyl, silyl, and acyl protecting groups or groups that possess substantially the same properties as these groups can be used.

An alkyl protecting group, as used herein, means triphenylmethyl or an alkyl group that possesses substantially the same protecting properties as triphenylmethyl. A silyl protecting group, as used herein, means a trialkylsilyl group having the formula:



wherein  $R_1$ ,  $R_2$ , and  $R_3$  may be lower-alkyl, e.g., methyl, ethyl, butyl, and alkyl possessing 5 carbon atoms or less; or phenyl. Furthermore,  $R_1$  may be identical to  $R_2$ ;  $R_1$ ,  $R_2$ , and  $R_3$  may all be




identical. Examples of silyl protecting groups include, but are not limited to, trimethylsilyl and t-butyldiphenylsilyl.

An acyl group, as used herein to describe an acyl protecting group (as in 1) or to describe a carboxylate (as in 4), is a group having the formula:




wherein R' is a lower alkyl, e.g., methyl, ethyl, butyl, and alkyl possessing 5 carbon atoms or less; substituted lower alkyl wherein the alkyl bears one, two, or more simple substituents, including, but not limited to, amino, carboxyl, hydroxy, phenyl, lower-alkoxy, e.g., methoxy and ethoxy; phenyl; substituted phenyl wherein the phenyl bears one, two, or more simple substituents, including, but not limited to, lower alkyl, halo, e.g., chloro and bromo, sulfato, sulfonyloxy, carboxyl, carbonyl, lower-alkoxy, e.g., carbomethoxy and carbethoxy, amino, mono- and di- lower alkylamino, e.g., methylamino, amido, hydroxy, lower alkoxy, e.g., methoxy and ethoxy, lower-alkanoyloxy, e.g., acetoxy.

A silyated pyrimidine base, as used herein, means a compound having the formula:



wherein X is either a trialkylsilyloxy or a trialkylsilylamino group, Z is a trialkylsilyl group, and Y is further described below. A trialkylsilyl group, as used herein, means a group having the formula:



wherein R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> may be lower-alkyl, e.g., methyl, ethyl, butyl, and alkyl possessing 5 carbon atoms or less, or phenyl. Furthermore, R<sub>1</sub> may be identical to R<sub>2</sub>; R<sub>1</sub>, R<sub>2</sub>, and R<sub>3</sub> may all be identical. Examples of trialkylsilyl groups include, but are not limited to, trimethylsilyl and t-butyldiphenylsilyl.

The silyated pyrimidine base may be substituted with various Y substituents, including, but not limited to, hydrogen, methyl, halo, alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl, at position 5 of the silyated pyrimidine base (Y substituent in Fig. 1) to modify the

properties, such as transport properties or the rate of metabolism, of the BCH-189 analog.

Illustrative examples of the synthesis of BCH-189 or BCH-189 analogs according to the present invention are given in Figs. 2, 3, and 4 and the following descriptions.

Figure 2 shows the synthesis of BCH-189 starting with allyl alcohol 7. A NaH oil suspension (4.5 g, 60%, 110 mmol) was washed with THF twice (100 ml x 2) and the resulting solid suspended in THF (300 ml). The suspension was cooled to 0°C, allyl alcohol 7 (6.8 ml, 100 mmol) was added dropwise, and the mixture was stirred for 30 minutes at 0°C. *t*-Butyl-diphenylsilyl chloride (25.8 ml, 100.8 mmol) was added dropwise at 0°C and the reaction mixture was stirred for 1 hour at 0°C. The solution was quenched with water (100 ml), and extracted with diethyl ether (200 ml x 2). The combined extracts were washed with water, dried over  $MgSO_4$ , filtered, concentrated, and the residue distilled under vacuum (90-100°C at 0.5-0.6 mm Hg) to give a colorless liquid 8 (28 g., 94 mmol, 94%). ( $^1H$  NMR: 7.70-7.35 (10H, m, aromatic-H); 5.93 (1H, m,  $H_2$ ); 5.37 (1H, dt,  $H_1$ ) J=1.4 and 14.4 Hz; 5.07 (1H, dt,  $H_1$ ) J=1.4 and 8.7 Hz; 4.21 (2H, m,  $H_3$ ); 1.07 (9H, s, *t*-Bu))

The silyl allyl ether 8 (15.5 g, 52.3 mmol) was dissolved in  $CH_2Cl_2$  (400 ml), and ozonized at -78°C. Upon

completion of ozonolysis, DMS (15 ml, 204 mmol, 3.9 eq) was added at -78°C and the mixture was warmed to room temperature and stirred overnight. The solution was washed with water (100 ml x 2), dried over MgSO<sub>4</sub>, filtered, concentrated, and distilled under vacuum (100-110°C at 0.5-0.6 mm Hg) to give a colorless liquid **2** (15.0 g, 50.3 mmol, 96%). (<sup>1</sup>H NMR: 9.74 (1H, s, H-CO); 7.70-7.35 (10H, m, aromatic-H); 4.21 (2H, s, -CH<sub>2</sub>); 1.22 (9H, s, t-Bu))

Silylated glycoaldehyde **2** (15.0 g, 50.3 mmol) was dissolved in toluene (200 ml) and thioglycolic acid (3.50 ml, 50.3 mmol) was added all at once. The solution was refluxed for 2 hours while the resulting water was removed with a Dean-Stark trap. The solution was cooled to room temperature and washed with saturated NaHCO<sub>3</sub> solution and the aqueous washings were extracted with diethyl ether (200 ml x 2). The combined extracts were washed with water (100 ml x 2), dried over MgSO<sub>4</sub>, filtered, and concentrated to give a colorless oil **10** (16.5 g, 44.3 mmol, 88%), which gradually solidified under vacuum. Recrystallization from hexane afforded a white solid **10** (15.8 g, 84%). (<sup>1</sup>H NMR: 7.72-7.38 (10H, m, aromatic-H); 5.53 (1H, t, H<sub>2</sub>) J=2.7 Hz; 3.93 (1H, dd, -CH<sub>2</sub>O) J=9.3 Hz; 3.81 (1H, d, 1H<sub>4</sub>) J=13.8 Hz; 3.79 (1H, dd, -CH<sub>2</sub>O); 3.58 (1H, d, 1H<sub>4</sub>); 1.02 (9H, s, t-Bu))

2-(t-Butyl-diphenylsilyloxy)-methyl-5-oxo-1,2-oxathiolane **10** (5.0 g, 13.42 mmol) was dissolved in toluene (150 ml) and the solution was cooled to -78°C. Dibal-H solution (14

ml, 1.0 M in hexanes, 14 mmol) was added dropwise, while the inside temperature was kept below  $-70^{\circ}\text{C}$  all the time. After the completion of the addition, the mixture was stirred for 30 minutes at  $-78^{\circ}\text{C}$ . Acetic anhydride (5 ml, 53 mmol) was added and the mixture was warmed to room temperature and stirred overnight. Water (5 ml) was added to the mixture and the resulting mixture was stirred for 1 hour at room temperature. The mixture was diluted with diethyl ether (300 ml),  $\text{MgSO}_4$  (40 g) was added, and the mixture was stirred vigorously for 1 hour at room temperature. The mixture was filtered, concentrated, and the residue flash chromatographed with 20% EtOAc in hexanes to give a colorless liquid **11** (3.60 g, 8.64 mmol, 64%), which was a 6:1 mixture of anomers. ( $^1\text{H}$  NMR of the major isomer: 7.70-7.35 (10H, m, aromatic-H); 6.63 (1H, d,  $\text{H}_5$ )  $J=4.4$  Hz; 5.47 (1H, t,  $\text{H}_2$ ); 4.20-3.60 (2H, m,  $-\text{CH}_2\text{O}$ ); 3.27 (1H, dd,  $1\text{H}_4$ )  $J=4.4$  and 11.4 Hz; 3.09 (1H, d,  $1\text{H}_4$ )  $J=11.4$  Hz; 2.02 (3H, s,  $\text{CH}_3\text{CO}$ ); 1.05 (9H, s, t-Bu);  $^1\text{H}$  NMR of the minor isomer: 7.70-7.35 (10H, m, aromatic-H); 6.55 (1H, d,  $\text{H}_5$ )  $J=3.9$  Hz; 5.45 (1H, t,  $\text{H}_2$ ); 4.20-3.60 (2H, m,  $-\text{CH}_2\text{O}$ ); 3.25 (1H, dd,  $1\text{H}_4$ )  $J=3.9$  and 11.4 Hz; 3.11 (1H, d,  $1\text{H}_4$ )  $J=11.4$  Hz; 2.04 (3H, s,  $\text{CH}_3\text{CO}$ ); 1.04 (9H, s, t-Bu))

2-(t-Butyl-diphenylsilyloxy)-methyl-5-acetoxy-1,3-oxathiolane **11** (0.28 g, 0.67 mmol) was dissolved in 1,2-dichloroethane (20 ml), and silylated cytosine **12** (0.20 g, 0.78 mmol) was added at once at room temperature. The mixture was stirred for 10 minutes and to it was added  $\text{SnCl}_4$  solution (0.80

ml, 1.0 M solution in  $\text{CH}_2\text{Cl}_2$ , 0.80 mmol) dropwise at room temperature. Additional cytosine **12** (0.10 g, 0.39 mmol) and  $\text{SnCl}_4$  solution (0.60 ml) were added in a same manner 1 hour later. After completion of the reaction in 2 hours, the solution was concentrated, and the residue was triturated with triethylamine (2 ml) and subjected to flash chromatography (first with neat EtOAc and then 20% ethanol in EtOAc) to give a tan solid **13** (100%  $\beta$  configuration) (0.25 g, 0.54 mmol, 80%). ( $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ): 7.75 (1H, d,  $\text{H}_6$ )  $J=7.5$  Hz; 7.65-7.35 (10H, m, aromatic-H); 7.21 and 7.14 (2H, broad,  $-\text{NH}_2$ ); 6.19 (1H, t,  $\text{H}_5$ ); 5.57 (1H, d,  $\text{H}_3$ ); 5.25 (1H, t,  $\text{H}_2$ ); 3.97 (1H, dd,  $-\text{CH}_2\text{O}$ )  $J=3.9$  and 11.1 Hz; 3.87 (1H, dd,  $-\text{CH}_2\text{O}$ ); 3.41 (1H, dd,  $1\text{H}_4$ )  $J=4.5$  and 11.7 Hz; 3.03 (1H, dd,  $1\text{H}_4$ )  $J=?$ ; 0.97 (9H, s, t-Bu))

Silyl ether **11** (0.23 g, 0.49 mmol) was dissolved in THF (30 ml), and to it was added  $n\text{-Bu}_4\text{NF}$  solution (0.50 ml, 1.0 M solution in THF, 0.50 mmol) dropwise at room temperature. The mixture was stirred for 1 hour and concentrated under vacuum. The residue was taken up with ethanol/triethylamine (2 ml/1 ml), and subjected to flash chromatography (first with EtOAc, then 20% ethanol in EtOAc) to afford a white solid **14** in 100% anomeric purity (BCH-189; 0.11 g, 0.48 mmol, 93%), which was further recrystallized from ethanol/ $\text{CHCl}_3$ /hexanes mixture. ( $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ ): 7.91 (1H, d,  $\text{H}_6$ )  $J=7.6$  Hz; 7.76 and 7.45 (2H, broad,  $-\text{NH}_2$ ); 6.19 (1H, t,  $\text{H}_5$ ); 5.80 (1H, d,  $\text{H}_3$ )  $J=7.6$  Hz; 5.34 (1H,

broad, -OH); 5.17 (1H, t, H<sub>2</sub>); 3.74 (2H, m, -CH<sub>2</sub>O); 3.42 (1H, dd, 1H<sub>4</sub>,) J=5.6 and 11.5 Hz; 3.09 (1H, dd, 1H<sub>4</sub>,) J=4.5 and 11.5 Hz)

BCH-189 and its analogs can also be synthesized by coupling a silylated uracil derivative with 11. Silylated uracil derivative 15 (1.80 g, 7.02 mmol) was coupled with 11 (1.72 g, 4.13 mmol) in 1,2-dichloroethane (50 ml) in the presence of SnCl<sub>4</sub> (5.0 ml) as described above in the the preparation of the cytsine derivative 13. The reaction was complete after 5 hours. Flash chromatography, first with 40% EtOAc in hexane and then EtOAc, afforded a white foam 16 (1.60 g, 3.43 mmol, 83%). (<sup>1</sup>H NMR: 9.39 (1H, broad, -NH) 7.90 (1H, d, H<sub>6</sub>) J=7.9 Hz; 7.75-7.35 (10H, m, aromatic-H); 6.33 (1H, dd, H<sub>5</sub>,); 5.51 (1H, d, H<sub>3</sub>) J=7.9 Hz; 5.23 (1H, t, H<sub>2</sub>); 4.11 (1H, dd, -CH<sub>2</sub>O) J=3.2 and 11.7 Hz; 3.93 (1H, dd, -CH<sub>2</sub>O); 3.48 (1H, dd, 1H<sub>4</sub>,) J=5.4 and 12.2 Hz; 3.13 (1H, dd, 1H<sub>4</sub>,) J=3.2 and 12.2 Hz)

The uracil derivative 16 can be converted to the cytosine derivative 13. The uracil derivative 16 (0.20 g, 0.43 mmol) was dissolved in a mixture of pyridine/dichloroethane (2 ml/10 ml), and the solution cooled to 0°C. Triflic anhydride (72 µl, 0.43 mmol) was added dropwise at 0°C and the mixture was warmed to room temperature and stirred for 1 hour. Additional triflic anhydride (0.50 µl, 0.30 mmol) was added and the mixture stirred for 1 hour. TLC showed no mobility with EtOAc. The reaction mixture was then decannulated into a NH<sub>3</sub>-saturated

methanol solution (30 ml) and the mixture was stirred for 12 hours at room temperature. The solution was concentrated, and the residue subjected to flash chromatography to give a tanned foam 11 (0.18 g, 0.39 mmol, 91%), which was identical with the compound obtained from the cytosine coupling reaction.

Fig. 3 illustrates the synthesis of 5-methylcytidine and thymidine derivatives of BCH-189. The acetate 11 (0.93 g, 2.23 mmol) in 1,2-dichloroethane (50 ml), was reacted with the silylated thymine derivative 17 (1.0 g, 3.70 mmol), and SnCl<sub>4</sub> solution (4.0 ml) in a manner similar to that described for the preparation of cytosine derivative 11. (<sup>1</sup>H NMR: 8.10 (1H, broad, NH); 7.75-7.30 (11H, m, 10 Aromatic H's and 1H<sub>6</sub>); 6.32 (1H, t, H<sub>1</sub>,) J=5.4 Hz; 5.25 (1H, t, H<sub>4</sub>,) J=4.2 Hz; 4.01 (1H, dd, 1H<sub>5</sub>,) J=3.9 and 11.4 Hz; 3.93 (1H, dd, 1H<sub>5</sub>,) J=4.5 and 11.4 Hz; 3.41 (1H, dd, 1H<sub>2</sub>,) J=5.4 and 11.7 Hz; 3.04 (1H, dd, 1H<sub>2</sub>,) J=5.7 and 11.7 Hz; 1.75 (3H, s, CH<sub>3</sub>); 1.07 (9H, s, t-Bu))

The thymine derivative 18 (0.20 g, 0.42 mmol) was dissolved in a mixture of pyridine/dichloroethane (2 ml/10 ml), and the solution cooled to 0°C. To it was added triflic anhydride (100 µl, 0.60 mmol) dropwise at 0°C, and the mixture was allowed, with continuous stirring, to warm to room temperature. After reaching room temperature, it was stirred for 1 hour. TLC showed no mobility with EtOAc. The reaction mixture was then decannulated into the NH<sub>3</sub>-saturated methanol solution



(20 ml), and the mixture stirred for 12 hours at room temperature. The solution was concentrated, and the residue was subjected to flash chromatography to give a tanned foam 19 (0.18 g, 0.38 mmol, 90%). (<sup>1</sup>H NMR: 7.70-7.30 (12H, m, 10 Aromatic H's, 1NH and H<sub>6</sub>); 6.60 (1H, broad, 1NH); 6.34 (1H, t, H<sub>1</sub>) J=4.5 Hz; 5.25 (1H, t, H<sub>4</sub>) J=3.6 Hz; 4.08 (1H, dd, 1H<sub>5</sub>) J=3.6 and 11.4 Hz; 3.96 (1H, dd, 1H<sub>5</sub>) J=3.6 and 11.4 Hz; 3.52 (1H, dd, 1H<sub>2</sub>) J=5.4 and 12.3 Hz; 3.09 (1H, dd, 1H<sub>2</sub>) J=3.9 and 12.3 Hz; 1.72 (3H, s, CH<sub>3</sub>); 1.07 (9H, s, t-Bu))

Silylother 19 (0.18 g, 0.38 mmol) was dissolved in THF (20 ml), and an n-Bu<sub>4</sub>NF solution (0.50 ml, 1.0 M solution in THF, 0.50 mmol) was added, dropwise, at room temperature. The mixture was stirred for 1 hour and concentrated under vacuum. The residue was taken up with ethanol/triethylamine (2 ml/1 ml), and subjected to flash chromatography (first with EtOAc, then 20% ethanol in EtOAc) to afford a white solid 20 (0.09 g, 0.37 mmol, 97%), which was further recrystallized from ethanol/CHCl<sub>3</sub>/Hexanes mixture to afford 82 mg of pure compound (89%). (<sup>1</sup>H NMR: (in d<sup>6</sup>-DMSO): 7.70 (1H, s, H<sub>6</sub>); 7.48 and 7.10 (2H, broad, NH<sub>2</sub>); 6.19 (1H, t, H<sub>1</sub>) J=6.5 Hz; 5.31 (1H, t, OH); 5.16 (1H, t, 1H<sub>4</sub>) J=5.4 Hz; 3.72 (2H, m, 2H<sub>5</sub>) 3.36 (1H, dd, 1H<sub>2</sub>) J=6.5 and 14.0 Hz; 3.05 (1H, dd, 1H<sub>2</sub>) J=6.5 and 14.0 Hz; 1.85 (3H, s, CH<sub>3</sub>))

Silylother 18 (0.70 g, 1.46 mmol) was dissolved in THF (50 ml), and an n-Bu<sub>4</sub>NF solution (2 ml, 1.0 M solution in THF, 2

mmol) was added, dropwise, at room temperature. The mixture was stirred for 1 hour and concentrated under vacuum. The residue was taken up with ethanol/triethylamine (2 ml/1 ml), and subjected to flash chromatography to afford a white solid **21** (0.33 g, 1.35 mmol, 92%). (<sup>1</sup>H NMR: (in d<sup>6</sup>-Acetone): 9.98 (1H, broad, NH); 7.76 (1H, d, H<sub>6</sub>) J=1.2 Hz; 6.25 (1H, t, H<sub>4</sub>) J=5.7 Hz; 5.24 (1H, t, H<sub>1</sub>) J=4.2 Hz; 4.39 (1H, t, OH) J=5.7 Hz; 3.85 (1H, dd, 2H<sub>3</sub>) J=4.2 and 5.7 Hz; 3.41 (1H, dd, 1H<sub>2</sub>) J=5.7 and 12.0 Hz; 3.19 (1H, dd, 1H<sub>2</sub>) J=5.4 and 12.0 Hz; 1.80 (3H, s, CH<sub>3</sub>))

Fig. 4 illustrates the synthesis of enantiomerically-enriched BCH-189 and its analogs. Allyl butyrate **22** (19.0 g, 148 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (400 ml), and ozonized at -78°C. Upon completion of ozonolysis, dimethyl sulfide (20 ml, 270 mmol, 1.8 eq) was added at -78°C and the mixture was warmed to room temperature and stirred overnight. The solution was washed with water (100 ml x 2), dried over MgSO<sub>4</sub>, filtered, concentrated, and distilled under vacuum (70-80°C at 0.5-0.6 mm Hg) to give a colorless liquid **21** (17.0 g, 131 mmol, 88%). (<sup>1</sup>H NMR: 9.59 (1H, s, H-CO); 4.66 (2H, s, -CH<sub>2</sub>O); 2.42 (2H, t, CH<sub>2</sub>CO) J=7.2 Hz; 1.71 (2H, sex, -CH<sub>2</sub>); 0.97 (3H, t, CH<sub>3</sub>) J=7.2 Hz) (IR (neat): 2990, 2960, 2900, 1750, 1740, 1460, 1420, 1390, 1280, 1190, 1110, 1060, 1020, 990, 880, 800, 760)

Butyryloxyacetaldehyde **21** (15.0 g, 115 mmol) was dissolved in toluene (200 ml) and mixed with thioglycolic acid

(8.0 ml, 115 mmol). The solution was refluxed for 5 hours while the resulting water was removed with a Dean-Stark trap. The solution was cooled to room temperature and was transferred to a 500 ml separatory funnel. The solution was then washed with saturated  $\text{NaHCO}_3$  solution. These aqueous washings were extracted with diethyl ether (200 ml x 2) to recuperate any crude product from the aqueous layer. The ether extracts were added to the toluene layer and the resulting mixture was washed with water (100 ml x 2), dried over  $\text{MgSO}_4$ , filtered, concentrated, and distilled under vacuum (70-80°C at 0.5-0.6 mm Hg) to give a colorless oil 24 (19 g, 93 mmol, 81%). ( $^1\text{H}$  NMR: 5.65 (1H, dd,  $\text{H}_3$ )  $J=5.0$  and 1.4 Hz; 4.35 (1H, dd,  $-\text{CH}_2\text{O}$ )  $J=3.2$  and 12.2 Hz; 4.29 (1H, dd,  $-\text{CH}_2\text{O}$ )  $J=5.7$  and 12.2 Hz; 3.72 (1H, d,  $-\text{CH}_2\text{S}$ )  $J=16.2$  Hz; 3.64 (1H, d,  $-\text{CH}_2\text{S}$ ; 2.34 (2H, t,  $-\text{CH}_2\text{CO}$ )  $J=7.2$  Hz; 1.66 (2H, sex,  $-\text{CH}_2$ ); 0.95 (3H, t,  $\text{CH}_3$ )  $J=7.2$  Hz) (IR (neat): 2980, 2960, 2900, 1780, 1740, 1460, 1410, 1390, 1350, 1300, 1290, 1260, 1220, 1170, 1110, 1080, 1070, 1000, 950, 910, 830, 820, 800, 760).

Pig liver esterase solution (90  $\mu\text{l}$ ) was added to a buffer solution (pH 7, 100 ml) at room temperature, and the mixture stirred vigorously for 5 minutes. The butyrate 24 (2.8 g, 13.7 mmol) was added, all at once, to the esterase/buffer solution and the mixture was stirred vigorously at room temperature for 2 hours. The reaction mixture was poured into a separatory funnel. The reaction flask was washed with ether (10

ml) and the washing was combined with the reaction mixture in the funnel. The combined mixture was extracted with hexanes three times (100 ml x 3). The three hexane extracts were combined and dried over  $\text{MgSO}_4$ , filtered, and concentrated to give the optically active butyrate 24 (1.12 g, 5.48 mmol, 40%). Enantiomeric excess was determined by an NMR experiment using a Tris(3-heptafluoropropyl-hydroxymethylene)-(+)-camphorato] europium (III) derivative as a chemical shift reagent; this procedure showed approximately 40% enrichment for one enantiomer. The remaining aqueous layer from the reaction was subjected to a continuous extraction with  $\text{CH}_2\text{Cl}_2$  for 20 hours. The organic layer was removed from the extraction apparatus, dried over  $\text{MgSO}_4$ , filtered, and concentrated to give an oil (1.24 g), which was shown by NMR analysis to consist of predominately the 2-hydroxymethyl-5-oxo-1,3-oxathiolane 25 with small amounts of butyric acid and the butyrate 24.

The lactone 25 (0.85 g, 4.16 mmol) was dissolved in toluene (30 ml), and the solution cooled to  $-78^\circ\text{C}$ . Dibal-H solution (9 ml, 1.0 M in hexanes, 9 mmol) was added dropwise, while the inside temperature was kept below  $-70^\circ\text{C}$  throughout the addition. After the addition was completed, the mixture was stirred for 0.5 hours at  $-78^\circ\text{C}$ . Acetic anhydride (5 ml, 53 mmol) was added and the mixture, with continuous stirring, was allowed to reach room temperature overnight. Water (5 ml) was added to the reaction mixture and the resultant mixture was stirred for 1

hour.  $\text{MgSO}_4$  (40 g) was then added and the mixture was stirred vigorously for 1 hour at room temperature. The mixture was filtered, concentrated, and the residue flash chromatographed with 20% EtOAc in hexanes to give a colorless liquid 26 (0.41 g, 1.86 mmol, 45%) which was a mixture of anomers at the C-4 position.

The 2-Acetoxymethyl-5-acetoxy-1,3-oxathiolane 26 (0.40 g, 1.82 mmol) was dissolved in 1,2-dichloroethane (40 ml), and to it the silylated cytosine 12 (0.70 g, 2.74 mmol) was added, all at once, at room temperature. The mixture was stirred for 10 minutes, and to it a  $\text{SnCl}_4$  solution (3.0 ml, 1.0 M solution in  $\text{CH}_2\text{Cl}_2$ , 3.0 mmol) was added, dropwise, at room temperature. Additional  $\text{SnCl}_4$  solution (1.0 ml) was added after 1 hour. The reaction was followed by TLC. Upon completion of the coupling, the solution was concentrated, the residue was triturated with triethylamine (2 ml) and subjected to flash chromatography (first with neat EtOAc then 20% ethanol in EtOAc) to give a tan solid 27 (0.42 g, 1.55 mmol, 86%). ( $^1\text{H}$  NMR: 7.73 (1H, d,  $\text{H}_6$ )  $J=7.5$  Hz; 6.33 (1H, t,  $\text{H}_4$ )  $J=4.8$  Hz; 5.80 (1H, d,  $\text{H}_5$ )  $J=7.5$  Hz; 4.52 (1H, dd,  $\text{H}_3$ )  $J=5.7$  and 12.3 Hz; 4.37 (1H, dd,  $\text{H}_3$ )  $J=3.3$  and 12.3 Hz; 3.54 (1H, dd,  $\text{H}_2$ )  $J=5.4$  and 12.0 Hz; 3.10 (1H, dd,  $\text{H}_3$ ); 2.11 (3H, s,  $\text{CH}_3$ ))


The 5'-Acetate of BCH-189 27 (140 mg, 0.52 mmol) was dissolved in anhydrous methanol (10 ml), and to it was added

sodium methoxide (110 mg, 2.0 mmol) in one portion. The mixture was stirred at room temperature until the hydrolysis was complete. The hydrolysis took about 1 hour, and the reaction was followed by TLC. Upon completion, the mixture was then concentrated, and the residue taken up with ethanol (2 ml). The ethanol solution was subjected to column chromatography using ethyl acetate first, then 20% ethanol in EtOAc to afford a white foam (110 mg, 92%), which exhibited an NMR spectrum identical to that of authentic BCH-189, 14.


WHAT IS CLAIMED IS:

1. A method of preparing the  $\beta$ -isomer of an antiviral nucleoside analog comprising the steps of:

(a) reducing a lactone having the formula:



wherein R is a protecting group, to form a carboxylate, said carboxylate having the formula:




wherein R' is an acyl group;

(b) coupling said carboxylate with a silylated pyrimidine base in the presence of an effective amount of  $\text{SnCl}_4$  to form the  $\beta$ -isomer of a 5' substituted 2',3'-dideoxy-3'-thia-nucleoside; and

(c) replacing said protecting group from the 5' position of said nucleoside with a hydrogen to form said antiviral nucleoside analog.

2. The method of Claim 1, wherein said protecting group is selected from the group consisting essentially of alkyl, silyl, and acyl.


3. The method of Claim 1, wherein said silyated pyrimidine base has the formula:



wherein X is selected from the group consisting essentially of trialkylsilyloxy and trialkylsilylamino;  
 wherein Y is selected from the group consisting essentially of hydrogen, methyl, halo, alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl; and  
 wherein Z is a trialkylsilyl group.

4. The method of Claim 1, wherein said antiviral nucleoside analog is BCH-189.

5. The method of Claim 1, wherein said antiviral nucleoside analog comprises the formula:




wherein Y is selected from the group consisting essentially of halo, alkyl, alkenyl, alkynyl, hydroxyalkyl,




carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.

6. The method of Claim 1, wherein said antiviral nucleoside analog comprises the formula:




wherein Y is selected from the group consisting essentially of hydrogen, halo, alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.

7. The method of Claim 1, wherein said antiviral nucleoside analog comprises the formula:



8. The method of Claim 1, wherein said antiviral nucleoside analog comprises the formula:



9. The method of Claim 1, further comprising the steps prior to (a) of:

(1) ozonizing a compound having the formula  $\text{CH}_2\text{CHCH}_2\text{OR}$  to form a glycoaldehyde having the formula  $\text{OHCCCH}_2\text{OR}$ , wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

(2) adding an effective amount of thioglycolic acid to said glycoaldehyde to form said lactone.

10. The method of Claim 1, wherein said reduction of said lactone is accomplished by addition of a reducing agent followed by addition of an effective amount of a carboxylic anhydride.


11. The method of Claim 10, wherein said reducing agent is selected from the group consisting essentially of DIBAL-H, RED-AL, and  $\text{NaBH}_4$ .

12. The method of Claim 1, wherein said replacement of said protecting group is accomplished by addition of an effective amount of  $(n\text{-C}_4\text{H}_9)_4\text{NF}$ .

13. The method of Claim 1, wherein said replacement of said protecting group is accomplished by addition of an effective amount of sodium methoxide.

14. A method of preparing an enantiomerically-enriched  $\beta$ -isomer of an antiviral nucleoside analog comprising the steps of:

(a) adding an effective amount of a stereoselective enzyme to a lactone having the formula:



wherein R is an acyl protecting group, to form enantiomerically-enriched 2-hydroxymethyl-5-oxo-1,3-oxathialane;


(b) reducing said enantiomerically-enriched 2-hydroxymethyl-5-oxo-1,3-oxathiolane to form an enantiomerically-enriched 2-acyloxymethyl-5-acyloxy-1,3-oxathiolane;

(c) coupling said enantiomerically-enriched 2-acyloxymethyl-5-acyloxy-1,3-oxathiolane with a silylated pyrimidine base in the presence of an effective amount of  $\text{SnCl}_4$  to form the  $\beta$ -isomer of a 2',3'-dideoxy-5'-acyloxymethyl-3'-thia-nucleoside; and

(d) replacing the 5'-acyloxymethyl substituent of said

nucleoside with a hydroxymethyl substituent to form said antiviral nucleoside analog.

15. The method of Claim 14, wherein said silyated pyrimidine base has the formula:




wherein X is selected from the group consisting essentially of trialkylsilyloxy and trialkylsilylamino;

wherein Y is selected from the group consisting essentially of hydrogen, methyl, halo, alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl; and

wherein Z is a trialkylsilyl group.


16. The method of Claim 14, wherein said antiviral nucleoside analog is BCH-189.

17. The method of Claim 14, wherein said antiviral nucleoside analog comprises the formula:




wherein Y is selected from the group consisting essentially of halo, alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.

18. The method of Claim 14, wherein said antiviral nucleoside analog comprises the formula:




wherein Y is selected from the group consisting essentially of hydrogen, halo, alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.

19. The method of Claim 14, wherein said antiviral nucleoside analog comprises the formula:



20. The method of Claim 14, wherein said antiviral nucleoside analog comprises the formula:



21. The method of Claim 14, further comprising the steps prior to (a) of:

(1) ozonizing a compound having the formula  $\text{CH}_2\text{CHCH}_2\text{OR}$  to form a glycoaldehyde having the formula  $\text{OHCCH}_2\text{OR}$ , wherein R is is an acyl group; and

(2) adding an effective amount of thioglycolic acid to said glycoaldehyde to form said lactone.


22. The method of Claim 14, wherein said reduction of said lactone is accomplished by addition of a reducing agent followed by addition of an effective amount of a carboxylic anhydride.

23. The method of Claim 22, wherein said reducing agent is selected from the group consisting essentially of DIBAL-H, RED-AL, and  $\text{NaBH}_4$ .

24. The method of Claim 14, wherein said stereoselective enzyme is selected from the group consisting of pig liver osterase, porcine pancreatic lipase, and subtilisin.

25. The method of Claim 14, wherein said replacement of said protecting group is accomplished by addition of an effective amount of sodium methoxide.

26. A method of preparing a carboxylate having the formula:




wherein R is a protecting group; and

wherein R' is an acyl group, comprising the steps of:

(a) ozonizing a compound having the formula  $\text{CH}_2\text{CHCH}_2\text{OR}$  to form a glycoaldehyde having the formula  $\text{OHCCH}_2\text{OR}$ , wherein R is a protecting group;

(b) adding an effective amount of thioglycolic acid to said glycoaldehyde to form a lactone having the formula:



; and


(c) reducing said lactone to form said carboxylate.

27. The method of Claim 26, wherein said reduction of said lactone is accomplished by a addition of a reducing agent followed by addition of an effective amount of a carboxylic anhydride.

28. The method of Claim 26, wherein said reducing agent is selected from the group consisting essentially of DIBAL-H, RED-AL, and  $\text{NaBH}_4$ .

29. The method of Claim 26, wherein said protecting group is selected from the group consisting essentially of alkyl, silyl, and acyl.


30. A carboxylate having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein R' is an acyl group.

31. An acetate having the formula:






wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl.

32. A method of preparing enantiomerically-enriched 2-acyloxymethyl-5-acyloxy-1,3-oxathiolane comprising the steps of:

(a) ozonizing a compound having the formula  $\text{CH}_2\text{CHCH}_2\text{OR}$  to form a glycoaldehyde having the formula  $\text{OHCCH}_2\text{OR}$ , wherein R is a protecting group;

(b) adding an effective amount of thioglycolic acid to said glycoaldehyde to form a lactone having the formula:



; and

(c) adding an effective amount of a stereoselective enzyme to said lactone to form enantiomerically-enriched 2-hydroxymethyl-5-oxo-1,3-oxathiolane; and

(d) reducing said enantiomerically-enriched 2-hydroxymethyl-5-oxo-1,3-oxathiolane to form enantiomerically-enriched 2-acyloxymethyl-5-acyloxy-1,3-oxathiolane.

33. The method of Claim 32, wherein said reduction of said enantiomerically-enriched 2-butyryloxymethyl-5-oxo-1,3-oxathiolane is accomplished by a addition of a reducing agent

followed by addition of an effective amount of a carboxylic anhydride.

34. The method of Claim 32, wherein said reducing agent is selected from the group consisting essentially of DIBAL-H, RED-AL, and  $\text{NaBH}_4$ .

35. The method of Claim 32, wherein said stereoselective enzyme is selected from the group consisting of pig liver osterase, porcine pancreatic lipase, and subtilisin.


36. The method of Claim 32, wherein said protecting group is selected from the group consisting essentially of alkyl, silyl, and acyl.

37. Enantiomerically-enriched 2-hydroxymethyl-5-oxo-1,3-oxathiolane.

38. Enantiomerically-enriched 2-acyloxymethyl-5-acyloxy-1,3-oxathiolane.

39. Enantiomerically-enriched 2-acetoxymethyl-5-acetoxy-1,3-oxathiolane.

40. A method of preparing the  $\beta$ -isomer of a substituted nucleoside comprising the step coupling a carboxylate having the formula:




wherein R is a protecting group; and

wherein R' is an acyl group, with a silylated pyrimidine base in the presence of an effective amount of  $\text{SnCl}_4$  to form the  $\beta$ -isomer of a 5' substituted 2',3'-dideoxy-3'-thia- nucleoside.

41. The method of Claim 40, wherein said protecting group is selected from the group consisting essentially of alkyl, silyl, and acyl.

42. The method of Claim 40, wherein said silylated pyrimidine base has the formula:




wherein X is selected from the group consisting essentially of trialkylsilyloxy and trialkylsilylamino;

wherein Y is selected from the group consisting essentially of hydrogen, methyl, halo, alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl; and

wherein Z is a trialkylsilyl group.


43. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is selected from the group consisting essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.


44. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is selected from the group consisting essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.


45. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is a hydrogen.


46. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is a hydrogen.


47. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is selected from the group consisting of chloro, bromo, fluoro, and iodo.


48. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is selected from the group consisting of chloro, bromo, fluoro, and iodo.


49. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is a methyl group.


50. A substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is a methyl group.


51. An enantiomerically-enriched substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is selected from the group consisting essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.


52. An enantiomerically-enriched substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is selected from the group consisting essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.


53. An enantiomerically-enriched substituted nucleoside having the formula:



wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and

wherein Y is a hydrogen.


54. An enantiomerically-enriched substituted nucleoside having the formula:






wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and  
 wherein Y is a hydrogen.

55. An enantiomerically-enriched substituted nucleoside having the formula:




wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and  
 wherein Y is selected from the group consisting essentially of chloro, bromo, fluoro, and iodo.

56. An enantiomerically-enriched substituted nucleoside having the formula:




wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and  
 wherein Y is selected from the group consisting essentially of chloro, bromo, fluoro, and iodo.

57. An enantiomerically-enriched substituted nucleoside having the formula:




wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and  
wherein Y is a methyl group.

58. An enantiomerically-enriched substituted nucleoside having the formula:




wherein R is selected from the group consisting essentially of alkyl, silyl, and acyl; and  
wherein Y is a methyl group.

59. A compound of the formula:




60. A compound of the formula:




wherein Y is selected from the group consisting of chloro, bromo, fluoro, and iodo.

61. A compound of the formula:




wherein Y is selected from the group consisting essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.

62. A compound of the formula:




63. A compound of the formula:




wherein Y is selected from the group consisting of chloro, bromo, fluoro, and iodo.

64. A compound of the formula:




wherein Y is selected from the group consisting essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl, carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl, cycloalkenyl, thioaryl, and selenoaryl.


65. A compound of the formula:



66. An enantiomerically-enriched compound of the formula:




67. An enantiomerically-enriched compound of the formula:




wherein Y is selected from the group consisting of  
chloro, bromo, fluoro, and iodo.

68. An enantiomerically-enriched compound of the formula:




wherein Y is selected from the group consisting  
essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl,  
carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl,  
cycloalkenyl, thioaryl, and selenoaryl.

69. An enantiomerically-enriched compound of the formula:




70. An enantiomerically-enriched compound of the formula:




wherein Y is selected from the group consisting of  
chloro, bromo, fluoro, and iodo.

71. An enantiomerically-enriched compound of the formula:



wherein Y is selected from the group consisting  
essentially of alkyl, alkenyl, alkynyl, hydroxyalkyl,  
carboxyalkyl, thioalkyl, selenoalkyl, phenyl, cycloalkyl,  
cycloalkenyl, thioaryl, and selenoaryl.

72. An enantiomerically-enriched compound of the formula:



# ABSTRACT

The present invention relates to a method of preparing BCH-189 and various analogs of BCH-189 from inexpensive precursors with the option of introducing functionality as needed. This synthetic route allows the stereoselective preparation of the biologically active isomer of these compounds,  $\beta$ -BCH-189 and related compounds. Furthermore, the stereochemistry at the nucleoside 4' position can be controlled to produce enantiomerically-enriched  $\beta$ -BCH-189 and its analogs.

c:\mpf\patent\emory\applicat.013\cek



EXPRESS MAIL NO. B011 781 05N

Docket No. 0510.013

DECLARATION FOR PATENT APPLICATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am an original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled "METHOD AND COMPOSITIONS FOR THE SYNTHESIS OF BCH-139 AND RELATED COMPOUNDS", the specification of which is attached hereto.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, §1.56(a).

I hereby appoint the following attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

William H. Needle (Reg. No. 26,209), Sumner C. Rosenberg (Reg. No. 28,753) and William G. Harvey (Reg. No. 32,573).

Address all telephone calls to William H. Needle at telephone no. (404) 688-0770.  
Address all correspondence to:

William H. Needle  
NEEDLE & ROSENBERG, P.C.  
133 Carnegie Way, N.W.  
Suite 400  
Atlanta, Georgia 30303

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first joint inventor: Dennis C. Liotta  
Inventor's signature: *[Signature]* Date: 02/01/90  
Residence: Stone Mountain, Georgia 30088 Citizenship: U.S.A.  
Post Office Address: 793 Post Road Way, Stone Mountain, Georgia 30088

Full name of second joint inventor: Woo-Bae Choi  
Inventor's signature: *[Signature]* Date: 02/01/90  
Residence: Atlanta, Georgia 30341 Citizenship: Republic of Korea  
Post Office Address: 3215A Flowers Road, S., Atlanta, Georgia 30341

c:\npf\forms\declarat\liotta.dec\cck

EXPRESS MAIL NO. B011 781 05N

07 473318

Page 1 of 2



Applicant or Patentee: Dennis C. Liotta et al. Attorney's  
Serial or Patent No.: Docket No.: 0510.013  
For: METHOD AND COMPOSITIONS FOR THE SYNTHESIS OF BCH-189 AND RELATED COMPOUNDS

**VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY  
STATUS (37 CFR 1.9(f) and 1.27(d)) - NONPROFIT ORGANIZATION**

I hereby declare that I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION: EMORY UNIVERSITY  
ADDRESS OF ORGANIZATION: 303 B Dental School, Atlanta, Georgia 30322

**TYPE OF ORGANIZATION:**

- ☒ University or other institution of higher education  
☐ Would qualify as tax exempt under Internal Revenue Service Code (26 USC 501(a) and 501(c) (3)) if located in The United States of America  
☐ Would qualify as nonprofit scientific or educational under statute of state of The United States of America if located in The United States of America  
(Name of state) )  
(Citation of statute) )

I hereby declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9 (e) for purposes of paying reduced fees under section 41(a) and (b) of Title 35, United States Code with regard to the invention entitled METHOD AND COMPOSITIONS FOR THE SYNTHESIS OF BCH-189 AND RELATED COMPOUNDS by inventor(s) Dennis C. Liotta and Woo-Bae Choi described in the specification filed herewith.

I hereby declare that rights under contract or law have been conveyed to and remain with the nonprofit organization with regard to the above identified invention.

If the rights held by the nonprofit organization are not exclusive, each individual, concern or organization having rights to the invention is listed below\* and no rights to the invention are held by any person, other than the inventor, who could not qualify as a small business concern under 37 CFR 1.9(d) or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

\*NOTE: Separate verified statements are required for each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

NAME \_\_\_\_\_  
ADDRESS \_\_\_\_\_  
☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING Ann R. Stevens, Ph.D.  
 TITLE IN ORGANIZATION Associate Vice President for Research  
 ADDRESS OF PERSON SIGNING Office of Sponsored Programs, EMORY UNIVERSITY,  
101 B Dental School, Atlanta, Georgia 30322  
 SIGNATURE Ann R. Stevens DATE 1/19/90

c:\upf\patent\emory\sm-ent11.013\cck


07 473318

Figure 1




473318

Figure 2




07 473318

Figure 3



01 473318

Figure 4



BCH-189

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:



☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**